



Lexicon Elects Diane E. Sullivan to Board of Directors

July 28, 2023

Ms. Sullivan brings her extensive commercialization, strategy, and market access experience to the Lexicon Board of Directors

The Woodlands, Texas, July 28, 2023 – Lexicon Pharmaceuticals, Inc. (Nasdaq: LXRX) today announced that Diane E. Sullivan has been elected to its Board of Directors.

Ms. Sullivan brings extensive commercialization, strategy, and market access experience to the Lexicon Board of Directors. She was chief commercial officer of The Medicines Company until its acquisition by Novartis, and subsequently served as chief commercial officer at DalCorp Pharmaceuticals. Prior to her time at The Medicines Company, Ms. Sullivan held senior positions at AstraZeneca, Pfizer, Wyeth, and GlaxoSmithKline in a variety of disciplines, including market access, strategy, marketing, brand management, business development and integration.

"I am looking forward to contributing to the company's important mission of pioneering medicines that transform patients' lives and helping to provide strategic direction to the company's commercial launch of INPEFA™ (sotagliflozin) and advancement of its pipeline," said Ms. Sullivan.

Ms. Sullivan served as a director of OrthogenRx, a privately held medical device company, until it was acquired by Avanos Medical in January 2022 and currently serves as a director of Amarin. Ms. Sullivan received her B.A. from Dickinson College.

Ms. Sullivan's appointment is concurrent with the resignation from the Lexicon Board of Directors of Frank Palantoni after 18 years of service culminating in the approval of INPEFA.

"We are grateful to Frank for his service on the Board of Directors and appreciate his contributions to the company's progress," said Ray Debbane, Lexicon's Board Chair. "With Diane's appointment, I am confident that the extensive pharmaceutical commercial experience she brings will provide Lexicon with valuable insight and direction as we pursue our launch of INPEFA and gear up for the advancement of our LX9211 clinical program. We are very excited to have Diane join the Board of Directors."

About Lexicon Pharmaceuticals

Lexicon is a biopharmaceutical company with a mission of pioneering medicines that transform patients' lives. Through its Genome5000™ program, Lexicon scientists studied the role and function of nearly 5,000 genes and identified more than 100 protein targets with significant therapeutic potential in a range of diseases. Through the precise targeting of these proteins, Lexicon is pioneering the discovery and development of innovative medicines to treat diseases safely and effectively. Lexicon is commercially launching one of these medicines, INPEFA™ (sotagliflozin), in the United States and has a pipeline of other promising drug candidates in discovery and preclinical and clinical development in neuropathic pain, diabetes and metabolism and other indications. For additional information, please visit www.lexpharma.com.

About INPEFA™ (sotagliflozin)

Discovered using Lexicon's unique approach to gene science, INPEFA™ (sotagliflozin) is an oral inhibitor of two proteins responsible for glucose regulation known as sodium-glucose cotransporter types 2 and 1 (SGLT2 and SGLT1). SGLT2 is responsible for glucose reabsorption by the kidney and SGLT1 is responsible for glucose absorption in the gastrointestinal tract. Sotagliflozin has been studied in multiple patient populations encompassing heart failure, diabetes, and chronic kidney disease in clinical studies involving approximately 20,000 patients.

INDICATION

INPEFA is indicated to reduce the risk of cardiovascular death, hospitalization for heart failure, and urgent heart failure visit in adults with:

- heart failure or
- type 2 diabetes mellitus, chronic kidney disease, and other cardiovascular risk factors

IMPORTANT SAFETY INFORMATION

Dosing: Assess renal function and volume status and, if necessary, correct volume depletion prior to initiation of INPEFA. INPEFA dosing for patients with decompensated heart failure may begin when patients are hemodynamically stable, including when

hospitalized or immediately upon discharge.

Contraindications: INPEFA is contraindicated in patients with hypersensitivity to any component.

Warnings and Precautions:

Ketoacidosis: INPEFA increases the risk of ketoacidosis in patients with type 1 diabetes mellitus (T1DM). Type 2 diabetes mellitus (T2DM) and pancreatic disorders are also risk factors. The risk of ketoacidosis may be greater with higher doses. There have been postmarketing reports of fatal events of ketoacidosis in patients with type 2 diabetes using sodium glucose transporter 2 (SGLT2) inhibitors. Before initiating INPEFA, assess risk factors for ketoacidosis. Consider ketone monitoring in patients with T1DM and consider ketone monitoring in others at risk for ketoacidosis, and educate patients on the signs/symptoms of ketoacidosis. Patients receiving INPEFA may require monitoring and temporary discontinuation of therapy in clinical situations known to predispose to ketoacidosis.

Assess patients who present with signs and symptoms of metabolic acidosis or ketoacidosis, regardless of blood glucose level. If suspected, discontinue INPEFA, evaluate, and treat promptly. Monitor patients for resolution of ketoacidosis before restarting INPEFA.

Volume Depletion: INPEFA can cause intravascular volume depletion which may sometimes manifest as symptomatic hypotension or acute transient changes in creatinine. There have been post-marketing reports of acute kidney injury, some requiring hospitalization and dialysis, in patients with type 2 diabetes mellitus receiving SGLT2 inhibitors. Patients with impaired renal function (eGFR < 60 mL/min/1.73 m²), elderly patients, or patients on loop diuretics may be at increased risk for volume depletion or hypotension. Before initiating INPEFA in patients with one or more of these characteristics, assess volume status and renal function, and monitor for signs and symptoms of hypotension during therapy.

Urosepsis and Pyelonephritis: Treatment with SGLT2 inhibitors, including INPEFA, increases the risk for urinary tract infections. Serious urinary tract infections including urosepsis and pyelonephritis requiring hospitalization have been reported. Evaluate patients for signs and symptoms of urinary tract infections and treat promptly.

Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues: Insulin and insulin secretagogues are known to cause hypoglycemia. INPEFA may increase the risk of hypoglycemia when combined with insulin or an insulin secretagogue. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used with INPEFA.

Necrotizing Fasciitis of the Perineum (Fournier's Gangrene): Reports of Fournier's Gangrene, a rare but serious and life-threatening necrotizing infection requiring urgent surgical intervention, have been identified in post-marketing surveillance in patients with diabetes mellitus receiving SGLT2 inhibitors. Assess patients who present with pain, tenderness, erythema, or swelling in the genital or perineal area, along with fever or malaise. If suspected, start treatment immediately with broad-spectrum antibiotics and, if necessary, surgical debridement. Discontinue INPEFA, closely monitor patient signs and symptoms, and provide appropriate alternative therapy for heart failure.

Genital Mycotic Infections: INPEFA increases the risk of genital mycotic infections. Monitor and treat as appropriate.

Urinary Glucose Test and 1,5-anhydroglucitol (1,5-AG) Assay: these are not reliable for patients taking SGLT2 inhibitors. Use alternative testing methods to monitor glucose levels.

Common Adverse Reactions: the most commonly reported adverse reactions (incidence ≥ 5%) were urinary tract infection, volume depletion, diarrhea, and hypoglycemia.

Drug Interactions:

- **Digoxin:** Monitor patients appropriately as there is an increase in the exposure of digoxin when coadministered with INPEFA 400 mg.
- **Uridine 5'-diphospho-glucuronosyltransferase (UGT) Inducer:** The coadministration of rifampicin, an inducer of UGTs, with sotagliflozin resulted in a decrease in the exposure of sotagliflozin.
- **Lithium:** Concomitant use of an SGLT2 inhibitor with lithium may decrease serum lithium concentrations. Monitor serum lithium concentration more frequently during INPEFA initiation and with dosage changes.

Use in Specific Populations:

- **Pregnancy and Lactation:** INPEFA is not recommended during the second and third trimesters of pregnancy, nor while breastfeeding.
- **Geriatric Use:** No INPEFA dosage change is recommended based on age. No overall differences in efficacy were detected between these patients and younger patients, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out. Elderly patients may be at increased risk for volume depletion adverse reactions, including hypotension.
- **Renal Impairment:** INPEFA was evaluated in patients with chronic kidney disease (eGFR 25 to 60 mL/min/1.73 m²) and in patients with heart failure with eGFR <60 mL/min/1.73 m². The safety profile of INPEFA across eGFR subgroups in these

studies was consistent with the known safety profile. There was an increase in volume-related adverse events (e.g., hypotension, dizziness) in patients with eGFR <30 mL/min/1.73m² relative to the overall safety population. Efficacy and safety studies with INPEFA did not enroll patients with an eGFR less than 25 mL/min/1.73 m² or on dialysis. After starting therapy in the studies, patients were discontinued if eGFR fell below 15 mL/min/1.73 m² or were initiated on chronic dialysis.

- **Hepatic Impairment:** INPEFA is not recommended in patients with moderate or severe hepatic impairment.

[Click here for full Prescribing Information.](#)

Safe Harbor Statement

This press release contains “forward-looking statements,” including statements relating to the therapeutic and commercial potential, research and clinical development and regulatory status of INPEFA™ (sotagliflozin), LX9211 and Lexicon’s other drug programs. In addition, this press release also contains forward looking statements relating to Lexicon’s financial position and long-term outlook on its business, growth and future operating results, discovery and development of products, strategic alliances and intellectual property, as well as other matters that are not historical facts or information. All forward-looking statements are based on management’s current assumptions and expectations and involve risks, uncertainties and other important factors, specifically including Lexicon’s ability to meet its capital requirements, successfully commercialize INPEFA in heart failure on the timeline and/or at the prices currently contemplated or at all, conduct preclinical and clinical development and obtain necessary regulatory approvals of INPEFA (in other indications), LX9211 and its other drug candidates on its anticipated timelines, achieve its operational objectives, obtain patent protection for its discoveries and establish strategic alliances, as well as additional factors relating to manufacturing, intellectual property rights, and the therapeutic or commercial value of its drug candidates. Any of these risks, uncertainties and other factors may cause Lexicon’s actual results to be materially different from any future results expressed or implied by such forward-looking statements. Information identifying such important factors is contained under “Risk Factors” in Lexicon’s annual report on Form 10-K for the year ended December 31, 2022 and other subsequent disclosure documents filed with the Securities and Exchange Commission. Lexicon undertakes no obligation to update or revise any such forward-looking statements, whether as a result of new information, future events or otherwise.

For Investor Inquiries:

Carrie Siragusa
Lexicon Pharmaceuticals, Inc.
csiragusa@lexpharma.com

For Media Inquiries:

Alina Cocuzza
Lexicon Pharmaceuticals, Inc.
acocuzza@lexpharma.com